



# From overnutrition to liver injury: AMP-activated protein kinase in nonalcoholic fatty liver diseases

Received for publication, May 28, 2020, and in revised form, June 24, 2020. Published, Papers in Press, July 10, 2020, DOI 10.1074/jbc.REV120.011356

Peng Zhao<sup>1</sup> and Alan R. Saltiel<sup>1,2,\*</sup>

From the Departments of <sup>1</sup>Medicine and <sup>2</sup>Pharmacology, University of California San Diego, La Jolla, California, USA

Edited by Qi-Qun Tang

Nonalcoholic fatty liver diseases (NAFLDs), especially nonalcoholic steatohepatitis (NASH), have become a major cause of liver transplant and liver-associated death. However, the pathogenesis of NASH is still unclear. Currently, there is no FDA-approved medication to treat this devastating disease. AMP-activated protein kinase (AMPK) senses energy status and regulates metabolic processes to maintain homeostasis. The activity of AMPK is regulated by the availability of nutrients, such as carbohydrates, lipids, and amino acids. AMPK activity is increased by nutrient deprivation and inhibited by overnutrition, inflammation, and hypersecretion of certain anabolic hormones, such as insulin, during obesity. The repression of hepatic AMPK activity permits the transition from simple steatosis to hepatocellular death; thus, activation might ameliorate multiple aspects of NASH. Here we review the pathogenesis of NAFLD and the impact of AMPK activity state on hepatic steatosis, inflammation, liver injury, and fibrosis during the transition of NAFL to NASH and liver failure.

Nonalcoholic fatty liver disease (NAFLD) is a major complication of metabolic dysfunction, usually a complication of obesity. NAFLD is an umbrella term describing two stages of chronic fatty liver diseases: NAFL (nonalcoholic fatty liver) and NASH (nonalcoholic steatohepatitis). NAFL is characterized by hepatic steatosis, which can be reversed simply by reduced caloric intake and exercise. Advanced stage NASH is characterized by steatosis with hepatic inflammation and liver injury, often accompanied by pericellular fibrosis (1). Although NASH is potentially reversible with diet and exercise, there is no FDA-approved medication to treat this devastating disease. NASH frequently progresses to cirrhosis, liver failure, and hepatocellular carcinoma (HCC) (1–3). The number of individuals with NAFLD in the United States is expected to increase from 83.1 million in 2015 to 100.9 million in 2030, whereas severe cases with advanced fibrosis are anticipated to increase by more than 100% (4).

Metabolic syndrome substantially increases the risk of NASH (5). The occurrence of NAFLD is predominantly associated with obesity and insulin resistance (6), whereas the incidence of NASH strongly correlates with central obesity, defined by the waist/hip ratio (7). Type 2 diabetes represents an independent pathogenic factor for NASH (8, 9). Most individuals with NAFLD exhibit dyslipidemia, including hypertriglyceridemia and hypercholesterolemia (8, 9). Moreover, diverticulosis and

overgrowth of the intestinal microbiome have been identified in human NASH. NASH-related liver injury and fibrosis might result from exposure to intestine-derived bacterial products, such as lipopolysaccharide (10). In addition, NAFLD and NASH can also result from a diverse array of pharmacotherapies, including glucocorticoids, tamoxifen, and methotrexate (8, 9) and long-term antiretroviral therapy for HIV (8).

The precise pathogenic mechanisms that give rise to NASH remain unclear. The “two-hit model” proposed that ectopic lipid storage caused by high-fat diet, obesity, and insulin resistance primes hepatocytes for a second insult inducing hepatic inflammation, liver injury, and fibrogenesis, which in turn promotes the progression from NAFL to NASH and cirrhosis (11, 12). However, it is now clear that multiple pathogenic factors may act in parallel and synergistically. A “multiple-hit model” proposes that multiple pathogenic factors act together to induce NAFLD, including but not limited to insulin resistance, inflammation, lipotoxicity, mitochondrial dysfunction, ER stress, oxidative stress, genetic determinants, and epigenetic factors (12).

AMP-activated protein kinase (AMPK) is an important energy sensor that regulates metabolic homeostasis. The activity of AMPK is inhibited by overnutrition during obesity and NAFLD (13–18). A recent study demonstrated that although the loss of AMPK activity does not affect hepatic lipid accumulation, it substantially exacerbates liver injury and hepatic fibrosis (17, 19), both of which could promote the transition from NASH to cirrhosis and HCC. Moreover, reactivation of AMPK improves symptoms of NASH and therapeutically improves liver injury (17, 19–21). We review here the regulation of AMPK activity during the pathophysiology of NAFL and the roles of the protein kinase in the regulation of NASH development.

## AMPK

AMPK is a heterotrimeric serine/threonine kinase comprised of three subunits: the  $\alpha$  catalytic subunit and the  $\beta$  and  $\gamma$  regulatory subunits. In mammalian cells, the AMPK  $\alpha$  subunit has two isoforms,  $\alpha 1$  and  $\alpha 2$ , encoded by the *Prkaa1* and *Prkaa2* genes. The  $\beta$  subunit has  $\beta 1$  and  $\beta 2$  isoforms, encoded by the *Prkab1* and *Prkab2* genes. The  $\gamma$  subunit includes three isoforms,  $\gamma 1$ ,  $\gamma 2$ , and  $\gamma 3$ , encoded by *Prkag1*, *Prkag2*, and *Prkag3*. These isoforms have the potential to form 12 different heterotrimeric complexes (15, 22). Although it remains uncertain whether there are functional differences among these isoforms, previous studies have reported different tissue distribution, regulation, subcellular localization, and functions for these complexes (23–25). The  $\gamma 2$  and  $\gamma 3$  isoforms possess

\*For correspondence: Alan R. Saltiel, [asaltiel@health.ucsd.edu](mailto:asaltiel@health.ucsd.edu).

unique N-terminal regions that may determine the subcellular localization of the holoenzyme (26).

The activity of AMPK is regulated by multiple factors through modulation of different subunits. The phosphorylation of Thr<sup>172</sup> within the catalytic domain of the  $\alpha$  subunit is required for the activation of AMPK (27, 28). Three upstream kinases, liver kinase B1 (LKB1),  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase kinase  $\beta$  (CaMKK $\beta$ ), and transforming growth factor  $\beta$  (TGF $\beta$ )-activated kinase 1 (TAK1), have been shown to phosphorylate residue Thr<sup>172</sup> (15, 29–34). Moreover, sirtuin 1 (SIRT1), an NAD<sup>+</sup>-dependent deacetylase, deacetylates LKB1 to induce its cytosolic localization and thus increases LKB1-dependent AMPK phosphorylation (35, 36). Another study showed that the Src family kinase Fyn phosphorylates LKB1 on Tyr<sup>265</sup> and Tyr<sup>365</sup> to increase LKB1 cytosolic localization and AMPK phosphorylation (37). Upon elevated intracellular  $\text{Ca}^{2+}$  concentrations, CaMKK $\beta$  directly phosphorylates AMPK on Thr<sup>172</sup> to increase its activity (29, 30, 33). Although the mechanism by which TAK1 activates AMPK is still unclear, several studies have demonstrated that the deletion of TAK1 inhibits starvation, metformin, and AICAR-induced AMPK activation (31, 34, 38).

As an important energy sensor, AMPK is allosterically activated by AMP (and ADP to a lesser extent), which binds to the  $\gamma$  subunit, and is repressed by ATP (22, 39). The interaction between AMP and the  $\gamma$  subunit leads to a conformational change, which protects Thr<sup>172</sup> from dephosphorylation (40–42). Consequently, AMPK senses a high AMP/ATP ratio and responds by increasing lipid oxidation and mitochondrial biogenesis, while reducing lipogenesis and glycogenesis, to increase intracellular ATP levels (43, 44). In addition to ATP, phosphocreatine also allosterically inhibits AMPK activity (45). This finding is consistent with the energy-sensing roles of AMPK.

AMPK activity is controlled by nutrients, including lipids, amino acids, and carbohydrates. High-fat diet feeding reduces AMPK expression and phosphorylation in skeletal muscle, heart, liver, adipose tissue, aortic endothelium, and hypothalamus (18, 46–52). One study suggested that palmitate represses AMPK activity via the ceramide-dependent activation of protein phosphatase 2A (PP2A). High-fat diets rich in palmitate inhibit AMPK activation *in vivo* (53). The increase of cardiac lipid content in Zucker rats and *ob/ob* mice results in the attenuation of AMPK activation. In cultured cardiomyocytes, fatty acids up-regulate the expression of *Ppm1a* (protein phosphatase 2C, PP2C) to inhibit AMPK, representing a feed-forward effect of lipid overload to promote energy storage (48). Excess amino acids have also been demonstrated to suppress AMPK activity (54, 55). High-protein diet or increased protein intake reduces AMPK phosphorylation while increasing mTOR phosphorylation in the hypothalamus and liver. An elevated level of amino acids, especially leucine, results in a decrease of the AMP/ATP ratio and thus represses AMPK activation (56, 57). The AMPK $\beta$  subunit also contains a conserved glycogen-binding domain. Glycogen, particularly when in a highly branched state, inhibits AMPK activity. Branched oligosaccharides with a single  $\alpha$ 1–6 branch allosterically inhibit AMPK phosphorylation by upstream kinases (58). Furthermore, recent studies demon-

strated that AMPK activity is modulated by glucose levels. Mechanistically, aldolase senses the glycolytic intermediate fructose 1,6-bisphosphate, and interacts with v-ATPase on the lysosomal surface. Without glucose, the absence of fructose 1,6-bisphosphate causes an altered interaction between aldolase and v-ATPase, which leads to the formation of an AMPK activation complex containing v-ATPase, LKB1, AMPK, AXIN, and Regulator, thus promoting AMPK $\alpha$  Thr<sup>172</sup> phosphorylation. The presence of glucose disrupts this complex and thus prevents AMPK activation (59).


Hormones and cytokines also modulate AMPK activity in physiological or pathological conditions. Insulin markedly reduces the activity of AMPK, both by increasing glucose uptake and oxidation and through Akt-mediated phosphorylation of AMPK $\alpha$  Ser<sup>485/491</sup>. These phosphorylation events inhibit the activity of the enzyme, leading to a conformational change that exposes the activating Thr<sup>172</sup> phosphorylation site within the kinase domain in the  $\alpha$  subunit. As a result, protein phosphatases, such as PP2A, dephosphorylate Thr<sup>172</sup> to deactivate AMPK (60). Proinflammatory cytokines, such as tumor necrosis factor  $\alpha$  (TNF $\alpha$ ), have also been shown to inhibit AMPK activity. TNF $\alpha$  induces the expression of *Ppm1a* (PP2C), which dephosphorylates Thr<sup>172</sup> to deactivate AMPK in skeletal muscle. Consequently, TNF $\alpha$  reduces ACC phosphorylation and represses fatty acid oxidation (14). Furthermore, TNF $\alpha$  activates TANK-binding kinase 1 (TBK1) in adipocytes. TBK1 phosphorylates Ser<sup>459</sup> and Ser<sup>476</sup> residues in the  $\alpha$  subunit to inhibit AMPK activity (18). Thus, AMPK activity is regulated by energy status, nutrient availability, and hormone/cytokine levels via various mechanisms, implicating its essential roles in monitoring metabolic processes (Fig. 1).

## AMPK and NAFLD

Several studies have reported a strong association between the reduction of AMPK activity and the incidence of metabolic diseases, including obesity, diabetes, and NAFLD (13). AMPK activity is inhibited in both obese rodents and human subjects, mainly attributable to the excess calorie intake and/or lack of exercise as well as increased inflammation (13–16, 18, 52). Hepatic AMPK activity is substantially attenuated in both NAFL and NASH (17, 19, 61). Although the roles of AMPK repression in the pathogenesis of these states remains uncertain, both pharmacological and genetic activation of AMPK in the liver exhibit beneficial effects on multiple aspects of NAFLD (17, 20, 62, 63).

### Hepatic steatosis

Ectopic lipid accumulation in the liver causes hepatic steatosis, which is tightly associated with obesity, insulin resistance, and diabetes. The amount of hepatic lipid content is predominantly regulated by four major pathways: *de novo* lipogenesis, fatty acid uptake, lipid oxidation, and very low-density lipoprotein (VLDL) secretion. *De novo* lipogenesis converts acetyl-CoA to fatty acids. The rate of *de novo* lipogenesis is determined by two key enzymes: acetyl-CoA carboxylase and fatty acid synthase (FAS). ACC catalyzes the carboxylation of acetyl-CoA to generate malonyl-CoA, whereas FAS uses acetyl-CoA



**Figure 1.** The regulation and function of AMPK

or malonyl-CoA to synthesize fatty acids. Esterification catalyzed by glycerol-3-phosphate acyltransferases, acylglycerol-3-phosphate acyltransferases, and diacylglycerol acyltransferases further converts fatty acids to triglycerides for storage (64). The increase of *de novo* lipogenesis and esterification results in hepatic steatosis. Activation of AMPK inhibits ACC via direct phosphorylation, reducing overall hepatic lipid storage, as seen in a phase 2 clinical trial (65). Hepatic lipid accumulation can also result from fatty acid uptake and subsequent esterification into triglyceride. Hepatic lipid accumulation increases as a function of high levels of serum-free fatty acids, which are determined by the rates of lipolysis in adipose tissue. In this regard, the ability of AMPK activation to improve systemic insulin sensitivity could indirectly lower lipolysis, thus reducing free fatty acid levels and fatty acid re-esterification in liver (66, 67).

*De novo* lipogenesis and fatty acid uptake are offset by lipid oxidation, the major catabolic pathway that resolves hepatic lipid storage.  $\beta$ -Oxidation converts fatty acids to acetyl-CoA. The rate-limiting step is catalyzed by carnitine palmitoyltransferases (CPTs), which transport cytosolic acyl-CoA into mitochondria. Acetyl-CoA generated from this process can enter the TCA cycle and then be utilized by mitochondria to produce ATP or heat. Thus, hepatic triglycerides and fatty acids levels are also tightly regulated by mitochondrial number and functions. Finally, hepatic lipid content is also regulated by VLDL packaging and secretion. VLDL is the major circulating vesicle that carries triglycerides from the liver to peripheral tissues (64).


The role of AMPK in hepatic lipid metabolism remains controversial. AMPK deletion does not dramatically affect hepatic steatosis under obesogenic conditions, although activation of AMPK by A-769662 reduced hepatic lipid content in high-fat-diet-induced NAFL (19). Mechanistically, AMPK inhibits *de novo* lipogenesis via directly phosphorylating ACC1 Ser<sup>79</sup> and ACC2 Ser<sup>212</sup> to repress the activity of the enzyme (19). The inhibition of ACC reduces malonyl-CoA, which is an allosteric inhibitor of CPT1 (68). Therefore, AMPK activation leads to increased fatty acid oxidation. In addition, sterol regulatory ele-

ment-binding proteins (SREBPs) regulate both triglyceride and cholesterol synthesis. Maturation and activation of SREBP-1c induce the expression of *Acc* and *Fasn* to increase *de novo* lipogenesis. Activation of SREBP-2 up-regulates the expression of cholesterol synthesis genes, including HMG-CoA reductase (*Hmgcr*), HMG-CoA synthase (*Hmgcs*), farnesyl diphosphate synthase (*Fdps*), and squalene synthase (*Sqs*) (69). AMPK directly phosphorylates and inhibits both SREBP-1 and -2 to reduce *de novo* lipogenesis and cholesterol synthesis in the liver (70). In addition, mTOR increases lipogenesis by promoting transcription and maturation of SREBPs (69). AMPK has been shown to phosphorylate both TSC2 and Raptor to inhibit mTORC1 activity (71–73). Consequently, activation of AMPK inhibits triglyceride and cholesterol synthesis to reduce hepatic steatosis.

The levels of circulating fatty acids are regulated by lipolysis in adipose tissue. Free fatty acids generated by adipocytes are burned through  $\beta$ -oxidation and respiration or secreted into the circulation. Enhanced lipolysis in adipose tissue increases circulating fatty acids, which in turn leads to hepatic fatty acid uptake and promotes steatosis (74, 75). On the other hand, increased fatty acid oxidation and mitochondrial respiration in adipose tissue reduces circulating fatty acids and alleviates hepatic steatosis (76). As a master regulator of metabolism, AMPK directly phosphorylates and activates peroxisome proliferator-activated receptor  $\gamma$  co-activator 1 $\alpha$  (PGC1 $\alpha$ ) to induce mitochondrial biogenesis, hence increasing mitochondrial number (77, 78). In addition, AMPK induces mitophagy via phosphorylating and activating Unc-51-like autophagy activating kinase 1 (ULK1) to clear damaged mitochondria and maintain mitochondrial homeostasis in adipose tissue (79, 80). Although it is still controversial, several studies indicated that AMPK directly phosphorylates hormone-sensitive lipase on Ser<sup>565</sup> to inhibit lipolysis in adipocytes (66, 81, 82). In summary, the activation of AMPK attenuates hepatic steatosis by modulating *de novo* lipogenesis, fatty acid oxidation, and fatty acid release from adipose tissue (Fig. 2).

### Hepatic inflammation

Inflammation is a hallmark for the progression from NAFL to NASH. The recruitment of immune cells, including macrophages, neutrophils, dendritic cells, and T cells, and the production of immune cell-derived cytokines, chemokines, and eicosanoids lead to hepatic inflammation (83). The increased number of recruited macrophages is currently used as a histological marker to determine liver inflammation. During the development of NASH, bone marrow-derived macrophages infiltrate into liver and work together with Kupffer cells to promote inflammation. Kupffer cells are yolk sac-derived, self-renewable liver-resident macrophages that localize within hepatic sinusoids (84). A recent study using single-cell RNA-Seq and lineage tracing has revealed that during NASH, resident Kupffer cell partially lose their cell identity and express genes that promote hepatocyte death. Meanwhile, increased bone marrow-derived macrophages acquire some Kupffer cell features and further enhance inflammation (85). Under inflammatory conditions during NASH, Kupffer cells can be induced to



**Figure 2. The regulation of hepatic steatosis by AMPK.** Activation of AMPK inhibits *de novo* lipogenesis while promoting fatty acid oxidation ( $\beta$ -oxidation) in the livers. In addition, AMPK activation reduces free fatty acid release from adipose tissue to prevent hepatic steatosis.

proliferate and differentiate into different subpopulations (86). Depending on the stimulating signals, both recruited macrophages and Kupffer cells undergo differentiation into M1- or M2-like macrophages (87). In response to proinflammatory stimuli, M1-like macrophages produce various cytokines, including TNF $\alpha$  and IL1 $\beta$ , to induce hepatocellular death and liver injury. In contrast, upon stimulation from signals inducing M2-like differentiation, macrophages secret cytokines like TGF $\beta$  to activate hepatic stellate cells (HSCs), and thus promote hepatic fibrosis (88–90). Therefore, hepatic inflammation plays a central role in the progression of NAFLD.

Oxidative stress, ER stress, lipotoxicity and mitochondrial dysfunction are among other pathogenic factors that trigger inflammation (83). Under conditions of oxidative stress, the production of reactive oxygen species (ROS) by mitochondria and NADPH oxidase is substantially up-regulated in the liver (91). The macrophage is a major source of ROS generated by NADPH oxidase (92, 93). Danger-associated molecular patterns are known to induce ROS formation in macrophages (94). ROS activates the NOD-, LRR-, and pyrin domain-containing protein-3 (NLRP3) inflammasome to induce inflammation. Blockage of the NLRP3 inflammasome attenuates hepatic inflammation and fibrosis in NASH (95). Oxidative stress promotes lipid peroxidation in the liver. The products of this non-enzymatic process, such as oxidized phospholipids and 4-hydroxynonenal, further enhance ROS generation to form a vicious cycle (96, 97). Neutralization of oxidized phospholipids prevents mitochondrial damage and protects against amylin diet-induced NASH (96). ER stress caused by the accumulation of unfolded or misfolded protein also promotes the production of ROS by inducing Ca $^{2+}$  release from the ER and thus inducing inflammation (98). Dietary factors, such as fructose, free fatty acids, and cholesterol, are external pathogenic factors triggering inflammation. Fructose is known to induce proinflammatory gene expression and impair  $\beta$ -oxidation in the liver (99–102). The amylin diet, containing fructose, is widely used to establish preclinical NASH mouse models. Elevated levels of

saturated fatty acids induce the accumulation of unfolded protein and ER stress (103). High free cholesterol has been associated with mitochondrial dysfunction, ER stress, and oxidative stress (104, 105), all of which contribute to the development of hepatic inflammation.

Previous studies demonstrated that activation of AMPK decreases the expression of proinflammatory mediators and attenuates inflammation in different conditions (106–108). Liver-specific expression of constitutively active AMPK reduces the expression of inflammatory genes (20). The chemokine CCL2 (monocyte chemoattractant protein 1, MCP-1) is an essential player in the recruitment of macrophages. The expression of *Ccl2* is induced by the activation of multiple proinflammatory signaling pathways, including NF $\kappa$ B- and JNK-mediated pathways (109, 110). AMPK activation by AICAR or by the expression of constitutively active AMPK largely alleviates palmitate- and TNF $\alpha$ -induced NF $\kappa$ B activation (111). Mechanistically, AMPK inhibits the nuclear localization of NF $\kappa$ B to repress the expression of NF $\kappa$ B target genes. Moreover, AMPK activation increases NAD $^+$  levels, leading to the activation of SIRT1. SIRT1 deacetylates the NF $\kappa$ B RelA/p65 subunit at Lys $^{310}$  to attenuate its transactivation activity (112). Our recent work found that AMPK phosphorylates ULK1 to induce TBK1 phosphorylation. TBK1, in turn, phosphorylates NF $\kappa$ B-inducing kinase (NIK) to induce its degradation. Consequently, AMPK activation leads to NIK degradation, resulting in attenuation of the atypical NF $\kappa$ B pathway, which is aberrantly activated in NAFLD (18, 113). Furthermore, AMPK activation by A-769662 inhibits IL-1 $\beta$ -induced JNK activation (114). As a result, AMPK inhibits proinflammatory signaling pathways to reduce *Ccl2* expression (18, 115). In addition, other studies have found that multiple downstream transcription factors, including FoxO family proteins and PGC1 $\alpha$ , could be involved in the anti-inflammatory effects of AMPK through regulating gene expression (108).

In addition to inhibiting proinflammatory signaling, AMPK may ameliorate inflammation via its anti-oxidative functions. ROS plays critical roles in the development of hepatic inflammation. AMPK activation attenuates cytosolic ROS production by down-regulating the expression of NAD(P)H oxidase genes and reducing mitochondrial ROS through an increase in PGC1 $\alpha$  target gene expression (116–119). Moreover, the activation of AMPK up-regulates the expression of *Sod2* (superoxide dismutase 2) and *Cat* (catalase) to alleviate oxidative stress (120). The reduction of ROS in turn ameliorates NLRP3 activation to reduce inflammation (121). Studies have demonstrated that antioxidants, such as coenzyme Q $_{10}$  and  $\gamma$ -tocotrienol, activate AMPK and inhibit NLRP3 activation (122, 123). An AMPK-FOXO3 pathway has been shown to induce the expression of thioredoxin (Trx) (124). Trx binds to the thioredoxin-interacting protein (Txnip), blocking the interaction between Txnip and NLRP3. Consequently, AMPK inhibits the activation of NLRP3 inflammasome to prevent hepatic inflammation (108, 125).

#### Liver injury

In normal liver, hepatocyte apoptosis maintains liver homeostasis, with a strict equilibrium between the loss and

replacement of hepatocytes (110, 126). However, under pathological conditions, such as viral infection, alcoholic or nonalcoholic steatohepatitis, and physical injury, extensive hepatocellular death leads to sustained liver injury, which is responsible for the enhanced scarring, bridging fibrosis, and subsequent development of cirrhosis (127–129). Moreover, hepatocellular death has been recognized as a major contributor to the progression to hepatocellular carcinoma (130). Therefore, understanding the molecular mechanisms of hepatocellular death is crucial for the treatment of liver diseases (131). Currently, hepatocellular death, reflected by increased serum aminotransferase levels, is the most widely used and sensitive parameter to screen for and monitor individuals with liver disease (110, 132). Evaluation of liver injury drives therapeutic decisions and has prognostic value for NASH.

Previous studies suggested that multiple types of cell death may contribute to liver injury in NASH. Although apoptosis plays a vital role to maintain homeostasis in healthy liver, elevated apoptotic stimuli produces extensive apoptosis, resulting in liver injury (126, 127). Whereas early findings suggested that the extensive apoptosis is responsible for NASH-associated liver damage, recent work suggests that other types of cell death also contribute to the pathogenesis of NASH. Necroptosis is a programmed form of inflammatory cell death that is mediated by the activation of receptor-interacting serine/threonine-protein kinase 3 (RIPK3). Activated RIPK3 phosphorylates mixed kinase domain-like protein, which forms pores in the membrane to cause rupture (133, 134). The expression of *Ripk3* is induced in a methionine- and choline-deficient diet (MCD)-induced NASH mouse model. Knockout of *Ripk3* ameliorates liver injury in these mice (135). Ferroptosis is a type of programmed cell death dependent on iron, producing lipid peroxidation-mediated cell death in NASH (136). Although it remains unclear whether pyroptosis, the highly inflammatory form of programmed cell death, occurs during NASH, the pyroptotic effector gasdermin D (GSDMD) and its pyroptosis-inducing fragment GSDMD-N are increased in human NASH. GSDMD deficiency alleviates lipogenesis and inflammation in MCD-induced NASH model (137).

In the pathogenesis of NASH, these pathogenic factors, including but not limited to oxidative stress, ER stress, lipotoxicity, and mitochondrial dysfunction, all activate signaling that has the capacity to mediate hepatocellular death (138). However, the crucial underlying mechanism for the regulation of hepatocellular death and liver injury during the transition from NAFL to NASH remains unclear. Our recent study found that normal AMPK activity is required to prevent hepatocellular death and liver damage. Liver-specific knockout of AMPK $\alpha 1/\alpha 2$  exaggerates liver injury in the choline-deficient high-fat diet (CD-HFD: 60% fat, 0.1% methionine, L-amino acid, no added choline)–induced NASH model. During the development of NASH, caspase-mediated apoptotic signaling pathways are induced in hepatocytes. The cleavage and activation of caspase-6 mediate a feed-forward loop to sustain the activation of apoptotic pathways and thus cause hepatocellular death in NASH. AMPK directly phosphorylates pro-caspase-6 to inhibit its cleavage and activation and control cell death. The repression


of AMPK during obesity and NAFLD unleashes caspase-6 to prime hepatocyte for apoptosis. We further demonstrated that activation of AMPK by A-769662 therapeutically improves liver damage even after NASH onset (17).

### Hepatic fibrosis

Hepatic fibrosis is a key feature used to determine severity of NASH. Progressive liver fibrosis frequently results in the progression from NASH to cirrhosis. Fibrosis is a wound-healing process that forms excess fibrous connective tissues to replace normal parenchymal tissues. Normal or mild fibrosis during injury is necessary for tissue repair. Extensive or chronic fibrosis results in the excessive accumulation of collagen and fibers in the extracellular space (62, 128). Hepatic fibrosis is induced upon activation of pathogenic pathways, including inflammation, oxidative stress, and liver injury. Whereas inflammation induces the pericellular fibrosis typically observed in the early stages of NASH, scarring after sustained liver injury leads to progressive fibrosis and subsequent development of cirrhosis (62, 128). The aggravation of liver damage worsens hepatic fibrosis even without changes in hepatic steatosis or inflammation (17). At advanced stages of hepatic fibrosis, the disruption of normal liver architecture and functions possibly results in the liver-associated death (128, 139). Therefore, resolving hepatic fibrosis and restoring liver functions are the ultimate goals for NASH treatment (1, 128, 139).

In chronic liver diseases, HSCs are direct mediators of fibrosis. Growth factors and inflammatory cytokines produced from other cell types, such as TGF $\beta$  from macrophages and platelet-derived growth factor from endothelial cells, cause overproliferation and transdifferentiation of HSCs. Upon activation, HSCs transdifferentiate into myofibroblasts, which produce an excessive amount of extracellular matrix (ECM) proteins (62, 140). Consequently, the accumulation of ECM and fibers disturbs hepatic homeostasis and further promotes the progression to cirrhosis or even HCC (141).

The activation of AMPK has been demonstrated to improve liver injury and attenuate hepatic fibrosis in different NASH models. AMPK activation by A-769662 improves liver injury and alleviates fibrosis in CD-HFD–induced NASH (17). Low doses of sorafenib, the first-line treatment for advanced HCC, activates AMPK and attenuates NASH-associated fibrosis in experimental mouse and monkey models (21). The injection of CCl<sub>4</sub> induces liver injury and hepatic fibrosis in a lipid-independent manner. In this mouse model, AMPK activation ameliorates fibrogenesis via inhibiting HSC proliferation and down-regulating the expression of fibrogenic genes, including *Nox4*, *Tgfb*, and *Acta2* (142, 143). Mechanistic studies further demonstrated that CCl<sub>4</sub> induces strong oxidative stress and hepatic accumulation of ROS, which in turn promotes hepatocellular death and liver fibrosis. The activation of AMPK prevents ROS production and HSCs activation and thus protects against liver injury and fibrosis (143–145). Furthermore, the fibrogenic cytokine TGF $\beta$  is mainly derived from macrophages and induces HSC activation during the fibrogenic process. Multiple studies demonstrated that the induction of AMPK activity represses TGF $\beta$ -induced expression of fibrogenic genes in



**Figure 3.** The effects of AMPK on hepatic steatosis, inflammation, liver injury, and fibrosis in NASH.

HSCs (62, 146–149). Both metformin and AICAR down-regulate the expression of *Col1a* and *Acta2* ( $\alpha$ -smooth muscle actin) in TGF $\beta$ -treated HSCs (150, 151). AMPK disrupts the interaction between Smad3 and its transcriptional coactivator p300 and induces proteasomal degradation of p300 to reduce fibrogenic genes expression in HSCs (151). Additionally, the activation of AMPK by adiponectin induces nitric oxide production to inhibit HSC proliferation and promote HSC apoptosis (149). AMPK activation by macrophage migration–inhibitory factor represses HSC migration and prevents hepatic fibrosis (62, 152). In summary, the activation of AMPK ameliorates hepatic fibrosis through multiple mechanisms, including reducing fibrogenic stimuli, preventing HSC activation/proliferation/migration and inhibiting expression of fibrogenic genes.

### Concluding remarks

AMPK is a critical energy sensor that regulates metabolic homeostasis. An increasing body of evidence has demonstrated that AMPK activity is repressed during metabolic disorders, including obesity, diabetes, and NAFLD. The inhibition of AMPK connects lipid dysregulation to inflammation, liver injury, and fibrosis in NAFLD (Fig. 3). Pharmacological activation of AMPK improves NASH in both murine and simian models (21). AMPK activators, such as A-769662, PF-739, or metformin, ameliorate symptoms of NASH-hepatic steatosis, inflammation, liver injury, and fibrosis via different mechanisms. However, global activation of AMPK by MK-8722 results in cardiomyocyte hypertrophy, possibly due to the induction of cardiac glycogen synthesis (153). Further investigation is needed to determine whether cardiomyocyte hypertrophy is a general effect for systemic AMPK activation or a side effect that is specific for MK-8722. In the event that increased AMPK activity results in cardiomyocyte hypertrophy, liver-specific activation of AMPK by liver cell–targeted drug delivery might be of great interest for the treatment of NASH.

R01DK117551, R01DK122804, and R01DK125820 (to A. R. S.) and National Institutes of Health Grant K99HL143277 (to P. Z.). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

**Conflict of interest**—A. R. S. and P. Z. are named inventors of patent application related to targeting the AMPK-caspase-6 axis for NASH treatment. A. R. S. is a founder of Elgia Therapeutics.

**Abbreviations**—The abbreviations used are: NAFLD, nonalcoholic fatty liver disease; NAFL, nonalcoholic fatty liver; NASH, nonalcoholic steatohepatitis; FDA, Food and Drug Administration; HCC, hepatocellular carcinoma; ER, endoplasmic reticulum; AMPK, AMP-activated protein kinase; TNF, tumor necrosis factor; VLDL, very low-density lipoprotein; FAS, fatty acid synthase; CPT, carnitine palmitoyltransferase; SREBP, sterol regulatory element–binding protein; mTOR, mechanistic target of rapamycin; HSC, hepatic stellate cell; ROS, reactive oxygen species; JNK, c-Jun N-terminal kinase; AICAR, 5-aminoimidazole-4-carboxamide ribonucleotide; NIK, NF $\kappa$ B-inducing kinase; MCD, methionine- and choline-deficient diet; GSDMD, gasdermin D; CD-HFD, choline-deficient high-fat diet; TGF, transforming growth factor; ECM, extracellular matrix.

### References

1. Friedman, S. L., Neuschwander-Tetri, B. A., Rinella, M., and Sanyal, A. J. (2018) Mechanisms of NAFLD development and therapeutic strategies. *Nat. Med.* **24**, 908–922 [CrossRef](#) [Medline](#)
2. Rinella, M. E., and Sanyal, A. J. (2016) Management of NAFLD: a stage-based approach. *Nat. Rev. Gastroenterol. Hepatol.* **13**, 196–205 [CrossRef](#) [Medline](#)
3. Lindenmeyer, C. C., and McCullough, A. J. (2018) The natural history of nonalcoholic fatty liver disease: an evolving view. *Clin. Liver Dis.* **22**, 11–21 [CrossRef](#) [Medline](#)
4. Estes, C., Razavi, H., Loomba, R., Younossi, Z., and Sanyal, A. J. (2018) Modeling the epidemic of nonalcoholic fatty liver disease demonstrates an exponential increase in burden of disease. *Hepatology* **67**, 123–133 [CrossRef](#) [Medline](#)
5. Marchesini, G., Bugianesi, E., Forlani, G., Cerrelli, F., Lenzi, M., Manini, R., Natale, S., Vanni, E., Villanova, N., Melchionda, N., and Rizzetto, M.

**Funding and additional information**—This work was supported by National Institutes of Health Grants P30DK063491, R01DK076906,

- (2003) Nonalcoholic fatty liver, steatohepatitis, and the metabolic syndrome. *Hepatology* **37**, 917–923 [CrossRef](#) [Medline](#)
6. Loomba, R., and Sanyal, A. J. (2013) The global NAFLD epidemic. *Nat. Rev. Gastroenterol. Hepatol.* **10**, 686–690 [CrossRef](#) [Medline](#)
  7. Farrell, G. C., and Larter, C. Z. (2006) Nonalcoholic fatty liver disease: from steatosis to cirrhosis. *Hepatology* **43**, S99–S112 [CrossRef](#) [Medline](#)
  8. Kopec, K. L., and Burns, D. (2011) Nonalcoholic fatty liver disease: a review of the spectrum of disease, diagnosis, and therapy. *Nutr. Clin. Pract.* **26**, 565–576 [CrossRef](#) [Medline](#)
  9. Reid, A. E. (2001) Nonalcoholic steatohepatitis. *Gastroenterology* **121**, 710–723 [CrossRef](#) [Medline](#)
  10. Shanab, A. A., Scully, P., Crosbie, O., Buckley, M., O'Mahony, L., Shanahan, F., Gazareen, S., Murphy, E., and Quigley, E. M. M. (2011) Small intestinal bacterial overgrowth in nonalcoholic steatohepatitis: association with toll-like receptor 4 expression and plasma levels of interleukin 8. *Dig. Dis. Sci.* **56**, 1524–1534 [CrossRef](#) [Medline](#)
  11. Peverill, W., Powell, L. W., and Skoien, R. (2014) Evolving concepts in the pathogenesis of NASH: beyond steatosis and inflammation. *Int. J. Mol. Sci.* **15**, 8591–8638 [CrossRef](#) [Medline](#)
  12. Buzzetti, E., Pinzani, M., and Tschochatzis, E. A. (2016) The multiple-hit pathogenesis of non-alcoholic fatty liver disease (NAFLD). *Metabolism* **65**, 1038–1048 [CrossRef](#) [Medline](#)
  13. Ruderman, N., and Prentki, M. (2004) AMP kinase and malonyl-CoA: targets for therapy of the metabolic syndrome. *Nat. Rev. Drug Discov.* **3**, 340–351 [CrossRef](#) [Medline](#)
  14. Steinberg, G. R., Michell, B. J., van Denderen, B. J., Watt, M. J., Carey, A. L., Fam, B. C., Andrikopoulos, S., Proietto, J., Görgün, C. Z., Carling, D., Hotamisligil, G. S., Febbraio, M. A., Kay, T. W., and Kemp, B. E. (2006) Tumor necrosis factor  $\alpha$ -induced skeletal muscle insulin resistance involves suppression of AMP-kinase signaling. *Cell Metab.* **4**, 465–474 [CrossRef](#) [Medline](#)
  15. Viollet, B., Horman, S., Leclerc, J., Lantier, L., Foretz, M., Billaud, M., Giri, S., and Andreelli, F. (2010) AMPK inhibition in health and disease. *Crit. Rev. Biochem. Mol. Biol.* **45**, 276–295 [CrossRef](#) [Medline](#)
  16. Xu, X. J., Gauthier, M. S., Hess, D. T., Apovian, C. M., Cacicedo, J. M., Gokce, N., Farb, M., Valentine, R. J., and Ruderman, N. B. (2012) Insulin sensitive and resistant obesity in humans: AMPK activity, oxidative stress, and depot-specific changes in gene expression in adipose tissue. *J. Lipid Res.* **53**, 792–801 [CrossRef](#) [Medline](#)
  17. Zhao, P., Sun, X., Chaggan, C., Liao, Z., In Wong, K., He, F., Singh, S., Loomba, R., Karin, M., Witztum, J. L., and Saltiel, A. R. (2020) An AMPK-caspase-6 axis controls liver damage in nonalcoholic steatohepatitis. *Science* **367**, 652–660 [CrossRef](#) [Medline](#)
  18. Zhao, P., Wong, K. I., Sun, X. L., Reilly, S. M., Uhm, M., Liao, Z. J., Skorobogatko, Y., and Saltiel, A. R. (2018) TBK1 at the crossroads of inflammation and energy homeostasis in adipose tissue. *Cell* **172**, 731–743 [CrossRef](#) [Medline](#)
  19. Boudaba, N., Marion, A., Huet, C., Pierre, R., Viollet, B., and Foretz, M. (2018) AMPK re-activation suppresses hepatic steatosis but its downregulation does not promote fatty liver development. *Ebiomedicine* **28**, 194–209 [CrossRef](#) [Medline](#)
  20. Garcia, D., Hellberg, K., Chaix, A., Wallace, M., Herzig, S., Badur, M. G., Lin, T., Shokhirev, M. N., Pinto, A. F. M., Ross, D. S., Saghatelian, A., Panda, S., Dow, L. E., Metallo, C. M., and Shaw, R. J. (2019) Genetic liver-specific AMPK activation protects against diet-induced obesity and NAFLD. *Cell Rep.* **26**, 192–208.e6 [CrossRef](#) [Medline](#)
  21. Jian, C., Fu, J., Cheng, X., Shen, L. J., Ji, Y. X., Wang, X., Pan, S., Tian, H., Tian, S., Liao, R., Song, K., Wang, H. P., Zhang, X., Wang, Y., Huang, Z., et al. (2020) Low-dose sorafenib acts as a mitochondrial uncoupler and ameliorates nonalcoholic steatohepatitis. *Cell Metab.* **31**, 892–908.e11 [CrossRef](#) [Medline](#)
  22. Miyaylova, M. M., and Shaw, R. J. (2011) The AMPK signalling pathway coordinates cell growth, autophagy and metabolism. *Nat. Cell Biol.* **13**, 1016–1023 [CrossRef](#) [Medline](#)
  23. Cheung, P. C., Salt, I. P., Davies, S. P., Hardie, D. G., and Carling, D. (2000) Characterization of AMP-activated protein kinase  $\gamma$ -subunit isoforms and their role in AMP binding. *Biochem. J.* **346**, 659–669 [CrossRef](#) [Medline](#)
  24. Cokorinos, E. C., Delmore, J., Reyes, A. R., Albuquerque, B., Kjobsted, R., Jorgensen, N. O., Tran, J. L., Jatkar, A., Cialdea, K., Esquejo, R. M., Meissen, J., Calabrese, M. F., Cordes, J., Moccia, R., Tess, D., et al. (2017) Activation of skeletal muscle AMPK promotes glucose disposal and glucose lowering in non-human primates and mice. *Cell Metab.* **25**, 1147–1159.e10 [CrossRef](#) [Medline](#)
  25. Salt, I., Celler, J. W., Hawley, S. A., Prescott, A., Woods, A., Carling, D., and Hardie, D. G. (1998) AMP-activated protein kinase: greater AMP dependence, and preferential nuclear localization, of complexes containing the  $\alpha 2$  isoform. *Biochem. J.* **334**, 177–187 [CrossRef](#) [Medline](#)
  26. Pinter, K., Grignani, R. T., Watkins, H., and Redwood, C. (2013) Localisation of AMPK  $\gamma$  subunits in cardiac and skeletal muscles. *J. Muscle Res. Cell Motil.* **34**, 369–378 [CrossRef](#) [Medline](#)
  27. Hawley, S. A., Davison, M., Woods, A., Davies, S. P., Beri, R. K., Carling, D., and Hardie, D. G. (1996) Characterization of the AMP-activated protein kinase kinase from rat liver and identification of threonine 172 as the major site at which it phosphorylates AMP-activated protein kinase. *J. Biol. Chem.* **271**, 27879–27887 [CrossRef](#) [Medline](#)
  28. Mitchelhill, K. I., Michell, B. J., House, C. M., Stapleton, D., Dyck, J., Gamble, J., Ullrich, C., Witters, L. A., and Kemp, B. E. (1997) Posttranslational modifications of the 5'-AMP-activated protein kinase  $\beta 1$  subunit. *J. Biol. Chem.* **272**, 24475–24479 [CrossRef](#) [Medline](#)
  29. Hawley, S. A., Pan, D. A., Mustard, K. J., Ross, L., Bain, J., Edelman, A. M., Frenguelli, B. G., and Hardie, D. G. (2005) Calmodulin-dependent protein kinase kinase- $\beta$  is an alternative upstream kinase for AMP-activated protein kinase. *Cell Metab.* **2**, 9–19 [CrossRef](#) [Medline](#)
  30. Hurley, R. L., Anderson, K. A., Franzone, J. M., Kemp, B. E., Means, A. R., and Witters, L. A. (2005) The  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinases are AMP-activated protein kinase kinases. *J. Biol. Chem.* **280**, 29060–29066 [CrossRef](#) [Medline](#)
  31. Momcilovic, M., Hong, S. P., and Carlson, M. (2006) Mammalian TAK1 activates Snf1 protein kinase in yeast and phosphorylates AMP-activated protein kinase *in vitro*. *J. Biol. Chem.* **281**, 25336–25343 [CrossRef](#) [Medline](#)
  32. Shaw, R. J., Kosmatka, M., Bardeesy, N., Hurley, R. L., Witters, L. A., DePinho, R. A., and Cantley, L. C. (2004) The tumor suppressor LKB1 kinase directly activates AMP-activated kinase and regulates apoptosis in response to energy stress. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 3329–3335 [CrossRef](#) [Medline](#)
  33. Woods, A., Dickerson, K., Heath, R., Hong, S. P., Momcilovic, M., Johnstone, S. R., Carlson, M., and Carling, D. (2005)  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase kinase-beta acts upstream of AMP-activated protein kinase in mammalian cells. *Cell Metab.* **2**, 21–33 [CrossRef](#) [Medline](#)
  34. Xie, M., Zhang, D., Dyck, J. R., Li, Y., Zhang, H., Morishima, M., Mann, D. L., Taffet, G. E., Baldini, A., Khouri, D. S., and Schneider, M. D. (2006) A pivotal role for endogenous TGF- $\beta$ -activated kinase-1 in the LKB1/AMP-activated protein kinase energy-sensor pathway. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 17378–17383 [CrossRef](#) [Medline](#)
  35. Lan, F., Cacicedo, J. M., Ruderman, N., and Ido, Y. (2008) SIRT1 modulation of the acetylation status, cytosolic localization, and activity of LKB1: possible role in AMP-activated protein kinase activation. *J. Biol. Chem.* **283**, 27628–27635 [CrossRef](#) [Medline](#)
  36. Howitz, K. T., Bitterman, K. J., Cohen, H. Y., Lamming, D. W., Lavu, S., Wood, J. G., Zipkin, R. E., Chung, P., Kisielewski, A., Zhang, L. L., Scherer, B., and Sinclair, D. A. (2003) Small molecule activators of sirtuins extend *Saccharomyces cerevisiae* lifespan. *Nature* **425**, 191–196 [CrossRef](#) [Medline](#)
  37. Yamada, E., Pessin, J. E., Kurland, I. J., Schwartz, G. J., and Bastie, C. C. (2010) Fyn-dependent regulation of energy expenditure and body weight is mediated by tyrosine phosphorylation of LKB1. *Cell Metab.* **11**, 113–124 [CrossRef](#) [Medline](#)
  38. Inokuchi-Shimizu, S., Park, E. J., Roh, Y. S., Yang, L., Zhang, B., Song, J., Liang, S., Pimienta, M., Taniguchi, K., Wu, X., Asahina, K., Lagakos, W., Mackey, M. R., Akira, S., Ellisman, M. H., et al. (2014) TAK1-mediated autophagy and fatty acid oxidation prevent hepatosteatosis and tumorigenesis. *J. Clin. Invest.* **124**, 3566–3578 [CrossRef](#) [Medline](#)

39. Hardie, D. G., Ross, F. A., and Hawley, S. A. (2012) AMPK: a nutrient and energy sensor that maintains energy homeostasis. *Nat. Rev. Mol. Cell Biol.* **13**, 251–262 [CrossRef](#) [Medline](#)
40. Suter, M., Riek, U., Tuerk, R., Schlattner, U., Wallimann, T., and Neumann, D. (2006) Dissecting the role of 5'-AMP for allosteric stimulation, activation, and deactivation of AMP-activated protein kinase. *J. Biol. Chem.* **281**, 32207–32216 [CrossRef](#) [Medline](#)
41. Sanders, M. J., Grondin, P. O., Hegarty, B. D., Snowden, M. A., and Carling, D. (2007) Investigating the mechanism for AMP activation of the AMP-activated protein kinase cascade. *Biochem. J.* **403**, 139–148 [CrossRef](#) [Medline](#)
42. Riek, U., Scholz, R., Konarev, P., Rufer, A., Suter, M., Nazabal, A., Ringler, P., Chami, M., Müller, S. A., Neumann, D., Forstner, M., Hennig, M., Zenobi, R., Engel, A., Svergun, D., et al. (2008) Structural properties of AMP-activated protein kinase: dimerization, molecular shape, and changes upon ligand binding. *J. Biol. Chem.* **283**, 18331–18343 [CrossRef](#) [Medline](#)
43. Daval, M., Foufelle, F., and Ferré, P. (2006) Functions of AMP-activated protein kinase in adipose tissue. *J. Physiol.* **574**, 55–62 [CrossRef](#) [Medline](#)
44. Hardie, D. G. (2011) AMP-activated protein kinase: an energy sensor that regulates all aspects of cell function. *Genes Dev.* **25**, 1895–1908 [CrossRef](#) [Medline](#)
45. Ponticos, M., Lu, Q. L., Morgan, J. E., Hardie, D. G., Partridge, T. A., and Carling, D. (1998) Dual regulation of the AMP-activated protein kinase provides a novel mechanism for the control of creatine kinase in skeletal muscle. *EMBO J.* **17**, 1688–1699 [CrossRef](#) [Medline](#)
46. Muse, E. D., Obici, S., Bhanot, S., Monia, B. P., McKay, R. A., Rajala, M. W., Scherer, P. E., and Rossetti, L. (2004) Role of resistin in diet-induced hepatic insulin resistance. *J. Clin. Invest.* **114**, 232–239 [CrossRef](#) [Medline](#)
47. Lee, W. J., Lee, I. K., Kim, H. S., Kim, Y. M., Koh, E. H., Won, J. C., Han, S. M., Kim, M. S., Jo, I., Oh, G. T., Park, I. S., Youn, J. H., Park, S. W., Lee, K. U., and Park, J. Y. (2005) α-Lipoic acid prevents endothelial dysfunction in obese rats via activation of AMP-activated protein kinase. *Arterioscler. Thromb. Vasc. Biol.* **25**, 2488–2494 [CrossRef](#) [Medline](#)
48. Wang, M. Y., and Unger, R. H. (2005) Role of PP2C in cardiac lipid accumulation in obese rodents and its prevention by troglitazone. *Am. J. Physiol. Endocrinol. Metab.* **288**, E216–E221 [CrossRef](#) [Medline](#)
49. Wilkes, J. J., Nguyen, M. T., Bandyopadhyay, G. K., Nelson, E., and Olefsky, J. M. (2005) Topiramate treatment causes skeletal muscle insulin sensitization and increased Acrp30 secretion in high-fat-fed male Wistar rats. *Am. J. Physiol. Endocrinol. Metab.* **289**, E1015–E1022 [CrossRef](#) [Medline](#)
50. Lessard, S. J., Chen, Z. P., Watt, M. J., Hashem, M., Reid, J. J., Febbraio, M. A., Kemp, B. E., and Hawley, J. A. (2006) Chronic rosiglitazone treatment restores AMPKα2 activity in insulin-resistant rat skeletal muscle. *Am. J. Physiol. Endocrinol. Metab.* **290**, E251–E257 [CrossRef](#) [Medline](#)
51. Liu, Y., Wan, Q., Guan, Q., Gao, L., and Zhao, J. (2006) High-fat diet feeding impairs both the expression and activity of AMPKα in rats' skeletal muscle. *Biochem. Biophys. Res. Commun.* **339**, 701–707 [CrossRef](#) [Medline](#)
52. Martin, T. L., Alquier, T., Asakura, K., Furukawa, N., Preitner, F., and Kahn, B. B. (2006) Diet-induced obesity alters AMP kinase activity in hypothalamus and skeletal muscle. *J. Biol. Chem.* **281**, 18933–18941 [CrossRef](#) [Medline](#)
53. Wu, Y., Song, P., Xu, J., Zhang, M., and Zou, M. H. (2007) Activation of protein phosphatase 2A by palmitate inhibits AMP-activated protein kinase. *J. Biol. Chem.* **282**, 9777–9788 [CrossRef](#) [Medline](#)
54. Leclerc, I., and Rutter, G. A. (2004) AMP-activated protein kinase: a new beta-cell glucose sensor?: regulation by amino acids and calcium ions. *Diabetes* **53**, S67–74 [CrossRef](#) [Medline](#)
55. Gleason, C. E., Lu, D., Witters, L. A., Newgard, C. B., and Birnbaum, M. J. (2007) The role of AMPK and mTOR in nutrient sensing in pancreatic beta-cells. *J. Biol. Chem.* **282**, 10341–10351 [CrossRef](#) [Medline](#)
56. Saha, A. K., Xu, X. J., Lawson, E., Deoliveira, R., Brandon, A. E., Kraegen, E. W., and Ruderman, N. B. (2010) Downregulation of AMPK accompanies leucine- and glucose-induced increases in protein synthesis and in-sulin resistance in rat skeletal muscle. *Diabetes* **59**, 2426–2434 [CrossRef](#) [Medline](#)
57. Chotechuang, N., Azzout-Marniche, D., Bos, C., Chaumontet, C., Gausseres, N., Steiler, T., Gaudichon, C., and Tome, D. (2009) mTOR, AMPK, and GCN2 coordinate the adaptation of hepatic energy metabolic pathways in response to protein intake in the rat. *Am. J. Physiol. Endocrinol. Metab.* **297**, E1313–E1323 [CrossRef](#) [Medline](#)
58. McBride, A., Ghilagaber, S., Nikolaev, A., and Hardie, D. G. (2009) The glycogen-binding domain on the AMPK β subunit allows the kinase to act as a glycogen sensor. *Cell Metab.* **9**, 23–34 [CrossRef](#) [Medline](#)
59. Lin, S. C., and Hardie, D. G. (2018) AMPK: sensing glucose as well as cellular energy status. *Cell Metab.* **27**, 299–313 [CrossRef](#) [Medline](#)
60. Suzuki, T., Bridges, D., Nakada, D., Skiniotis, G., Morrison, S. J., Lin, J. D., Saltiel, A. R., and Inoki, K. (2013) Inhibition of AMPK catabolic action by GSK3. *Mol. Cell* **50**, 407–419 [CrossRef](#) [Medline](#)
61. Qiang, X., Xu, L., Zhang, M., Zhang, P., Wang, Y., Wang, Y., Zhao, Z., Chen, H., Liu, X., and Zhang, Y. (2016) Demethyleneberberine attenuates non-alcoholic fatty liver disease with activation of AMPK and inhibition of oxidative stress. *Biochem. Biophys. Res. Commun.* **472**, 603–609 [CrossRef](#) [Medline](#)
62. Liang, Z., Li, T., Jiang, S., Xu, J., Di, W., Yang, Z., Hu, W., and Yang, Y. (2017) AMPK: a novel target for treating hepatic fibrosis. *Oncotarget* **8**, 62780–62792 [CrossRef](#) [Medline](#)
63. Smith, B. K., Marcinko, K., Desjardins, E. M., Lally, J. S., Ford, R. J., and Steinberg, G. R. (2016) Treatment of nonalcoholic fatty liver disease: role of AMPK. *Am. J. Physiol. Endocrinol. Metab.* **311**, E730–E740 [CrossRef](#) [Medline](#)
64. Nguyen, P., Leray, V., Diez, M., Serisier, S., Le Bloch, J., Siliart, B., and Dumon, H. (2008) Liver lipid metabolism. *J. Anim. Physiol. Anim. Nutr. (Berl.)* **92**, 272–283 [CrossRef](#) [Medline](#)
65. Alkhouri, N., Lawitz, E., Noureddin, M., DeFronzo, R., and Shulman, G. I. (2020) GS-0976 (Firsocostat): an investigational liver-directed acetyl-CoA carboxylase (ACC) inhibitor for the treatment of non-alcoholic steatohepatitis (NASH). *Expert Opin. Investig. Drugs* **29**, 135–141 [CrossRef](#) [Medline](#)
66. Daval, M., Diot-Dupuy, F., Bazin, R., Hainault, I., Viollet, B., Vaulont, S., Hajduch, E., Ferré, P., and Foufelle, F. (2005) Anti-lipolytic action of AMP-activated protein kinase in rodent adipocytes. *J. Biol. Chem.* **280**, 25250–25257 [CrossRef](#) [Medline](#)
67. Jeon, S. M. (2016) Regulation and function of AMPK in physiology and diseases. *Exp. Mol. Med.* **48**, e245 [CrossRef](#) [Medline](#)
68. McGarry, J. D., Takabayashi, Y., and Foster, D. W. (1978) The role of malonyl-CoA in the coordination of fatty acid synthesis and oxidation in isolated rat hepatocytes. *J. Biol. Chem.* **253**, 8294–8300 [Medline](#)
69. Han, J., and Wang, Y. (2018) mTORC1 signaling in hepatic lipid metabolism. *Protein Cell* **9**, 145–151 [CrossRef](#) [Medline](#)
70. Li, Y., Xu, S. Q., Mihaylova, M. M., Zheng, B., Hou, X. Y., Jiang, B. B., Park, O., Luo, Z. J., Lefai, E., Shyy, J. Y. J., Gao, B., Wierzbicki, M., Verbeuren, T. J., Shaw, R. J., Cohen, R. A., et al. (2011) AMPK phosphorylates and inhibits SREBP activity to attenuate hepatic steatosis and atherosclerosis in diet-induced insulin-resistant mice. *Cell Metab.* **13**, 376–388 [CrossRef](#) [Medline](#)
71. González, A., Hall, M. N., Lin, S. C., and Hardie, D. G. (2020) AMPK and TOR: the yin and yang of cellular nutrient sensing and growth control. *Cell Metab.* **31**, 472–492 [CrossRef](#) [Medline](#)
72. Inoki, K., Zhu, T., and Guan, K. L. (2003) TSC2 mediates cellular energy response to control cell growth and survival. *Cell* **115**, 577–590 [CrossRef](#) [Medline](#)
73. Gwinn, D. M., Shackelford, D. B., Egan, D. F., Mihaylova, M. M., Mery, A., Vasquez, D. S., Turk, B. E., and Shaw, R. J. (2008) AMPK phosphorylation of raptor mediates a metabolic checkpoint. *Mol. Cell* **30**, 214–226 [CrossRef](#) [Medline](#)
74. Thorne, A., Lofgren, P., and Hoffstedt, J. (2010) Increased visceral adipocyte lipolysis—a pathogenic role in nonalcoholic fatty liver disease? *J. Clin. Endocrinol. Metab.* **95**, E209–E213 [CrossRef](#)
75. Wueest, S., Item, F., Lucchini, F. C., Challa, T. D., Muller, W., Bluher, M., and Konrad, D. (2016) Mesenteric fat lipolysis mediates obesity-

- associated hepatic steatosis and insulin resistance. *Diabetes* **65**, 140–148 CrossRef Medline
76. Scheja, L., and Heeren, J. (2016) Metabolic interplay between white, beige, brown adipocytes and the liver. *J. Hepatol.* **64**, 1176–1186 CrossRef Medline
77. Cantó, C., and Auwerx, J. (2009) PGC-1 $\alpha$ , SIRT1 and AMPK, an energy sensing network that controls energy expenditure. *Curr. Opin. Lipidol.* **20**, 98–105 CrossRef Medline
78. Puigserver, P., Wu, Z., Park, C. W., Graves, R., Wright, M., and Spiegelman, B. M. (1998) A cold-inducible coactivator of nuclear receptors linked to adaptive thermogenesis. *Cell* **92**, 829–839 CrossRef Medline
79. Kim, J., Kundu, M., Viollet, B., and Guan, K. L. (2011) AMPK and mTOR regulate autophagy through direct phosphorylation of Ulk1. *Nat. Cell Biol.* **13**, 132–141 CrossRef Medline
80. Mottillo, E. P., Desjardins, E. M., Crane, J. D., Smith, B. K., Green, A. E., Ducommun, S., Henriksen, T. I., Rebalka, I. A., Razi, A., Sakamoto, K., Scheele, C., Kemp, B. E., Hawke, T. J., Ortega, J., Granneman, J. G., et al. (2016) Lack of adipocyte AMPK exacerbates insulin resistance and hepatic steatosis through brown and beige adipose tissue function. *Cell Metab.* **24**, 118–129 CrossRef Medline
81. Corton, J. M., Gillespie, J. G., Hawley, S. A., and Hardie, D. G. (1995) 5-Aminoimidazole-4-carboxamide ribonucleoside: a specific method for activating AMP-activated protein kinase in intact cells? *Eur. J. Biochem.* **229**, 558–565 CrossRef Medline
82. Sullivan, J. E., Brocklehurst, K. J., Marley, A. E., Carey, F., Carling, D., and Beri, R. K. (1994) Inhibition of lipolysis and lipogenesis in isolated rat adipocytes with AICAR, a cell-permeable activator of AMP-activated protein kinase. *FEBS Lett.* **353**, 33–36 CrossRef Medline
83. Schuster, S., Cabrera, D., Arrese, M., and Feldstein, A. E. (2018) Triggering and resolution of inflammation in NASH. *Nat. Rev. Gastroenterol. Hepatol.* **15**, 349–364 CrossRef Medline
84. Dixon, L. J., Barnes, M., Tang, H., Pritchard, M. T., and Nagy, L. E. (2013) Kupffer cells in the liver. *Compr. Physiol.* **3**, 785–797 CrossRef Medline
85. Seidman, J. S., Troutman, T. D., Sakai, M., Gola, A., Spann, N. J., Bennett, H., Bruni, C. M., Ouyang, Z., Li, R. Z., Sun, X., Vu, B. T., Pasillas, M. P., Ego, K. M., Gosselin, D., Link, V. M., et al. (2020) Niche-specific reprogramming of epigenetic landscapes drives myeloid cell diversity in nonalcoholic steatohepatitis. *Immunity* **52**, 1057–1074.e7 CrossRef Medline
86. Jenkins, S. J., Ruckerl, D., Cook, P. C., Jones, L. H., Finkelman, F. D., van Rooijen, N., MacDonald, A. S., and Allen, J. E. (2011) Local macrophage proliferation, rather than recruitment from the blood, is a signature of TH2 inflammation. *Science* **332**, 1284–1288 CrossRef Medline
87. Wan, J., Benkdane, M., Teixeira-Clerc, F., Bonnafous, S., Louvet, A., Lafidil, F., Pecker, F., Tran, A., Gual, P., Mallat, A., Lotersztajn, S., and Pavone, C. (2014) M2 Kupffer cells promote M1 Kupffer cell apoptosis: a protective mechanism against alcoholic and nonalcoholic fatty liver disease. *Hepatology* **59**, 130–142 CrossRef Medline
88. Zhang, F., Wang, H., Wang, X., Jiang, G., Liu, H., Zhang, G., Wang, H., Fang, R., Bu, X., Cai, S., and Du, J. (2016) TGF- $\beta$  induces M2-like macrophage polarization via SNAIL-mediated suppression of a pro-inflammatory phenotype. *Oncotarget* **7**, 52294–52306 CrossRef Medline
89. Mosser, D. M., and Edwards, J. P. (2008) Exploring the full spectrum of macrophage activation. *Nat. Rev. Immunol.* **8**, 958–969 CrossRef Medline
90. Koyama, Y., and Brenner, D. A. (2017) Liver inflammation and fibrosis. *J. Clin. Invest.* **127**, 55–64 CrossRef Medline
91. Masarone, M., Rosato, V., Dallio, M., Gravina, A. G., Aglitti, A., Loguerio, C., Federico, A., and Persico, M. (2018) Role of oxidative stress in pathophysiology of nonalcoholic fatty liver disease. *Oxid. Med. Cell Longev.* **2018**, 9547613 CrossRef Medline
92. Krenkel, O., and Tacke, F. (2017) Liver macrophages in tissue homeostasis and disease. *Nat. Rev. Immunol.* **17**, 306–321 CrossRef Medline
93. Liang, S., Ma, H. Y., Zhong, Z., Dhar, D., Liu, X., Xu, J., Koyama, Y., Nishio, T., Karin, D., Karin, G., McCubbin, R., Zhang, C., Hu, R., Yang, G., Chen, L., et al. (2019) NADPH oxidase 1 in liver macrophages promotes inflammation and tumor development in mice. *Gastroenterology* **156**, 1156–1172.e6 CrossRef Medline
94. Chatterjee, S., Rana, R., Corbett, J., Kadiiska, M. B., Goldstein, J., and Mason, R. P. (2012) P2X7 receptor-NADPH oxidase axis mediates protein radical formation and Kupffer cell activation in carbon tetrachloride-mediated steatohepatitis in obese mice. *Free Radic. Biol. Med.* **52**, 1666–1679 CrossRef Medline
95. Mridha, A. R., Wree, A., Robertson, A. A. B., Yeh, M. M., Johnson, C. D., Van Rooyen, D. M., Haczeyni, F., Teoh, N. C., Savard, C., Ioannou, G. N., Masters, S. L., Schroder, K., Cooper, M. A., Feldstein, A. E., and Farrell, G. C. (2017) NLRP3 inflammasome blockade reduces liver inflammation and fibrosis in experimental NASH in mice. *J. Hepatol.* **66**, 1037–1046 CrossRef Medline
96. Sun, X., Seidman, J. S., Zhao, P., Troutman, T. D., Spann, N. J., Que, X., Zhou, F., Liao, Z., Pasillas, M., Yang, X., Magida, J. A., Kisileva, T., Brenner, D. A., Downes, M., Evans, R. M., et al. (2020) Neutralization of oxidized phospholipids ameliorates non-alcoholic steatohepatitis. *Cell Metab.* **31**, 189–206.e8 CrossRef Medline
97. Dunning, S., Hannivoort, R. A., de Boer, J. F., Buist-Homan, M., Faber, K. N., and Moshage, H. (2009) Superoxide anions and hydrogen peroxide inhibit proliferation of activated rat stellate cells and induce different modes of cell death. *Liver Int.* **29**, 922–932 CrossRef Medline
98. Zhang, K., and Kaufman, R. J. (2008) From endoplasmic-reticulum stress to the inflammatory response. *Nature* **454**, 455–462 CrossRef Medline
99. Yang, Z. H., Miyahara, H., Takeo, J., and Katayama, M. (2012) Diet high in fat and sucrose induces rapid onset of obesity-related metabolic syndrome partly through rapid response of genes involved in lipogenesis, insulin signalling and inflammation in mice. *Diabetol. Metab. Syndr.* **4**, 32 CrossRef Medline
100. Liu, J., Zhuang, Z. J., Bian, D. X., Ma, X. J., Xun, Y. H., Yang, W. J., Luo, Y., Liu, Y. L., Jia, L., Wang, Y., Zhu, M. L., Ye, D. W., Zhou, G., Lou, G. Q., and Shi, J. P. (2014) Toll-like receptor-4 signalling in the progression of non-alcoholic fatty liver disease induced by high-fat and high-fructose diet in mice. *Clin. Exp. Pharmacol. Physiol.* **41**, 482–488 CrossRef Medline
101. Crescenzo, R., Bianco, F., Falcone, I., Coppola, P., Liverini, G., and Iossa, S. (2013) Increased hepatic *de novo* lipogenesis and mitochondrial efficiency in a model of obesity induced by diets rich in fructose. *Eur. J. Nutr.* **52**, 537–545 CrossRef Medline
102. Rebollo, A., Roglans, N., Baena, M., Sánchez, R. M., Merlos, M., Alegret, M., and Laguna, J. C. (2014) Liquid fructose downregulates Sirt1 expression and activity and impairs the oxidation of fatty acids in rat and human liver cells. *Biochim. Biophys. Acta* **1841**, 514–524 CrossRef Medline
103. Leamy, A. K., Egnatchik, R. A., and Young, J. D. (2013) Molecular mechanisms and the role of saturated fatty acids in the progression of non-alcoholic fatty liver disease. *Prog. Lipid Res.* **52**, 165–174 CrossRef Medline
104. Min, H. K., Kapoor, A., Fuchs, M., Mirshahi, F., Zhou, H., Maher, J., Kellum, J., Warnick, R., Contos, M. J., and Sanyal, A. J. (2012) Increased hepatic synthesis and dysregulation of cholesterol metabolism is associated with the severity of nonalcoholic fatty liver disease. *Cell Metab.* **15**, 665–674 CrossRef Medline
105. Al-Rasadi, K., Rizzo, M., Montalto, G., and Berg, G. (2015) Nonalcoholic fatty liver disease, cardiovascular risk, and carotid inflammation. *Angiology* **66**, 601–603 CrossRef Medline
106. Huang, B. P., Lin, C. H., Chen, H. M., Lin, J. T., Cheng, Y. F., and Kao, S. H. (2015) AMPK activation inhibits expression of proinflammatory mediators through downregulation of PI3K/p38 MAPK and NF- $\kappa$ B signaling in murine macrophages. *DNA Cell Biol.* **34**, 133–141 CrossRef Medline
107. O'Neill, L. A., and Hardie, D. G. (2013) Metabolism of inflammation limited by AMPK and pseudo-starvation. *Nature* **493**, 346–355 CrossRef Medline
108. Salminen, A., Hyttinen, J. M., and Kaarniranta, K. (2011) AMP-activated protein kinase inhibits NF- $\kappa$ B signaling and inflammation: impact on healthspan and lifespan. *J. Mol. Med. (Berl.)* **89**, 667–676 CrossRef Medline
109. Win, S., Than, T. A., Zhang, J., Oo, C., Min, R. W. M., and Kaplowitz, N. (2018) New insights into the role and mechanism of c-Jun-N-terminal kinase signaling in the pathobiology of liver diseases. *Hepatology* **67**, 2013–2024 CrossRef Medline

110. Luedde, T., and Schwabe, R. F. (2011) NF- $\kappa$ B in the liver—linking injury, fibrosis and hepatocellular carcinoma. *Nat. Rev. Gastroenterol. Hepatol.* **8**, 108–118 [CrossRef](#) [Medline](#)
111. Cacicedo, J. M., Yagihashi, N., Keaney, J. F., Jr., Ruderman, N. B., and Ido, Y. (2004) AMPK inhibits fatty acid-induced increases in NF- $\kappa$ B transactivation in cultured human umbilical vein endothelial cells. *Biochem. Biophys. Res. Commun.* **324**, 1204–1209 [CrossRef](#) [Medline](#)
112. Yeung, F., Hoberg, J. E., Ramsey, C. S., Keller, M. D., Jones, D. R., Frye, R. A., and Mayo, M. W. (2004) Modulation of NF- $\kappa$ B-dependent transcription and cell survival by the SIRT1 deacetylase. *EMBO J.* **23**, 2369–2380 [CrossRef](#) [Medline](#)
113. Xiong, Y., Torsoni, A. S., Wu, F., Shen, H., Liu, Y., Zhong, X., Canet, M. J., Shah, Y. M., Omary, M. B., Liu, Y., and Rui, L. (2018) Hepatic NF- $\kappa$ B-inducing kinase (NIK) suppresses mouse liver regeneration in acute and chronic liver diseases. *Elife* **7**, e34152 [CrossRef](#) [Medline](#)
114. Mancini, S. J., White, A. D., Bjelland, S., Rutherford, C., Graham, D., Richter, E. A., Viollet, B., Touyz, R. M., Palmer, T. M., and Salt, I. P. (2017) Activation of AMP-activated protein kinase rapidly suppresses multiple pro-inflammatory pathways in adipocytes including IL-1 receptor-associated kinase-4 phosphorylation. *Mol. Cell Endocrinol.* **440**, 44–56 [CrossRef](#) [Medline](#)
115. Peng, X., Li, J., Wang, M., Qu, K., and Zhu, H. (2019) A novel AMPK activator improves hepatic lipid metabolism and leukocyte trafficking in experimental hepatic steatosis. *J. Pharmacol. Sci.* **140**, 153–161 [CrossRef](#) [Medline](#)
116. Rabinovitch, R. C., Samborska, B., Faubert, B., Ma, E. H., Gravel, S. P., Andrzejewski, S., Raissi, T. C., Pause, A., St-Pierre, J., and Jones, R. G. (2017) AMPK maintains cellular metabolic homeostasis through regulation of mitochondrial reactive oxygen species. *Cell Rep.* **21**, 1–9 [CrossRef](#) [Medline](#)
117. Wang, S., Zhang, M., Liang, B., Xu, J., Xie, Z., Liu, C., Viollet, B., Yan, D., and Zou, M. H. (2010) AMPK $\alpha$ 2 deletion causes aberrant expression and activation of NAD(P)H oxidase and consequent endothelial dysfunction *in vivo*: role of 26S proteasomes. *Circ. Res.* **106**, 1117–1128 [CrossRef](#) [Medline](#)
118. Eid, A. A., Ford, B. M., Block, K., Kasinath, B. S., Gorin, Y., Ghosh-Choudhury, G., Barnes, J. L., and Abboud, H. E. (2010) AMP-activated protein kinase (AMPK) negatively regulates Nox4-dependent activation of p53 and epithelial cell apoptosis in diabetes. *J. Biol. Chem.* **285**, 37503–37512 [CrossRef](#) [Medline](#)
119. Song, P., and Zou, M. H. (2012) Regulation of NAD(P)H oxidases by AMPK in cardiovascular systems. *Free Radic. Biol. Med.* **52**, 1607–1619 [CrossRef](#) [Medline](#)
120. Garrido-Maraver, J., Paz, M. V., Cordero, M. D., Bautista-Lorite, J., Oropesa-Ávila, M., de la Mata, M., Pavón, A. D., de Lavera, I., Alcocer-Gómez, E., Galán, F., Ybot González, P., Cotán, D., Jackson, S., and Sánchez-Alcázar, J. A. (2015) Critical role of AMP-activated protein kinase in the balance between mitophagy and mitochondrial biogenesis in MELAS disease. *Biochim. Biophys. Acta* **1852**, 2535–2553 [CrossRef](#) [Medline](#)
121. Cordero, M. D., Williams, M. R., and Ryffel, B. (2018) AMP-activated protein kinase regulation of the NLRP3 inflammasome during aging. *Trends Endocrinol. Metab.* **29**, 8–17 [CrossRef](#) [Medline](#)
122. Bullón, P., Alcocer-Gómez, E., Carrión, A. M., Marín-Aguilar, F., Garrido-Maraver, J., Román-Malo, L., Ruiz-Cabello, J., Culic, O., Ryffel, B., Apetoh, L., Ghiringhelli, F., Battino, M., Sánchez-Alcázar, J. A., and Cordero, M. D. (2016) AMPK phosphorylation modulates pain by activation of NLRP3 inflammasome. *Antioxid. Redox Signal.* **24**, 157–170 [CrossRef](#) [Medline](#)
123. Kim, Y., Wang, W., Okla, M., Kang, I., Moreau, R., and Chung, S. (2016) Suppression of NLRP3 inflammasome by  $\gamma$ -tocotrienol ameliorates type 2 diabetes. *J. Lipid Res.* **57**, 66–76 [CrossRef](#) [Medline](#)
124. Li, X. N., Song, J., Zhang, L., LeMaire, S. A., Hou, X., Zhang, C., Coselli, J. S., Chen, L., Wang, X. L., Zhang, Y., and Shen, Y. H. (2009) Activation of the AMPK-FOXO3 pathway reduces fatty acid-induced increase in intracellular reactive oxygen species by upregulating thioredoxin. *Diabetes* **58**, 2246–2257 [CrossRef](#) [Medline](#)
125. Zhou, R., Tardivel, A., Thorens, B., Choi, I., and Tschoop, J. (2010) Thioredoxin-interacting protein links oxidative stress to inflammasome activation. *Nat. Immunol.* **11**, 136–140 [CrossRef](#) [Medline](#)
126. Michalopoulos, G. K., and DeFrances, M. (2005) Liver regeneration. *Adv. Biochem. Eng. Biotechnol.* **93**, 101–134 [CrossRef](#) [Medline](#)
127. Guicciardi, M. E., and Gores, G. J. (2005) Apoptosis: a mechanism of acute and chronic liver injury. *Gut* **54**, 1024–1033 [CrossRef](#) [Medline](#)
128. Hernandez-Gea, V., and Friedman, S. L. (2011) Pathogenesis of liver fibrosis. *Annu. Rev. Pathol.* **6**, 425–456 [CrossRef](#) [Medline](#)
129. Malhi, H., Guicciardi, M. E., and Gores, G. J. (2010) Hepatocyte death: a clear and present danger. *Physiol. Rev.* **90**, 1165–1194 [CrossRef](#) [Medline](#)
130. Feldstein, A. E., Canbay, A., Angulo, P., Taniai, M., Burgart, L. J., Lindor, K. D., and Gores, G. J. (2003) Hepatocyte apoptosis and Fas expression are prominent features of human nonalcoholic steatohepatitis. *Gastroenterology* **125**, 437–443 [CrossRef](#) [Medline](#)
131. Cao, L., Quan, X. B., Zeng, W. J., Yang, X. O., and Wang, M. J. (2016) Mechanism of hepatocyte apoptosis. *J. Cell Death* **9**, 19–29 [CrossRef](#) [Medline](#)
132. Hadizadeh, F., Faghihimani, E., and Adibi, P. (2017) Nonalcoholic fatty liver disease: diagnostic biomarkers. *World J. Gastrointest. Pathophysiol.* **8**, 11–26 [CrossRef](#) [Medline](#)
133. Dhuriya, Y. K., and Sharma, D. (2018) Necroptosis: a regulated inflammatory mode of cell death. *J. Neuroinflammation* **15**, 199 [CrossRef](#) [Medline](#)
134. Linkermann, A., and Green, D. R. (2014) Necroptosis. *N. Engl. J. Med.* **370**, 455–465 [CrossRef](#) [Medline](#)
135. Gautheron, J., Vucur, M., Reisinger, F., Cardenas, D. V., Roderburg, C., Koppe, C., Kreggenwinkel, K., Schneider, A. T., Bartneck, M., Neumann, U. P., Canbay, A., Reeves, H. L., Luedde, M., Tacke, F., Trautwein, C., et al. (2014) A positive feedback loop between RIP3 and JNK controls non-alcoholic steatohepatitis. *EMBO Mol. Med.* **6**, 1062–1074 [CrossRef](#) [Medline](#)
136. Qi, J., Kim, J. W., Zhou, Z., Lim, C. W., and Kim, B. (2020) Ferroptosis affects the progression of nonalcoholic steatohepatitis via the modulation of lipid peroxidation-mediated cell death in mice. *Am. J. Pathol.* **190**, 68–81 [CrossRef](#) [Medline](#)
137. Xu, B., Jiang, M., Chu, Y., Wang, W., Chen, D., Li, X., Zhang, Z., Zhang, D., Fan, D., Nie, Y., Shao, F., Wu, K., and Liang, J. (2018) Gasdermin D plays a key role as a pyroptosis executor of non-alcoholic steatohepatitis in humans and mice. *J. Hepatol.* **68**, 773–782 [CrossRef](#) [Medline](#)
138. Kanda, T., Matsuoka, S., Yamazaki, M., Shibata, T., Nirei, K., Takahashi, H., Kaneko, T., Fujisawa, M., Higuchi, T., Nakamura, H., Matsumoto, N., Yamagami, H., Ogawa, M., Imazu, H., Kuroda, K., et al. (2018) Apoptosis and non-alcoholic fatty liver diseases. *World J. Gastroenterol.* **24**, 2661–2672 [CrossRef](#) [Medline](#)
139. Wattacheril, J., Issa, D., and Sanyal, A. (2018) Nonalcoholic steatohepatitis (NASH) and hepatic fibrosis: emerging therapies. *Annu. Rev. Pharmacol. Toxicol.* **58**, 649–662 [CrossRef](#) [Medline](#)
140. Lee, U. E., and Friedman, S. L. (2011) Mechanisms of hepatic fibrogenesis. *Best Pract. Res. Clin. Gastroenterol.* **25**, 195–206 [CrossRef](#) [Medline](#)
141. Baglieri, J., Brenner, D. A., and Kisseleva, T. (2019) The role of fibrosis and liver-associated fibroblasts in the pathogenesis of hepatocellular carcinoma. *Int. J. Mol. Sci.* **20**, 1723 [CrossRef](#) [Medline](#)
142. Li, J., Pan, Y., Kan, M., Xiao, X., Wang, Y., Guan, F., Zhang, X., and Chen, L. (2014) Hepatoprotective effects of berberine on liver fibrosis via activation of AMP-activated protein kinase. *Life Sci.* **98**, 24–30 [CrossRef](#) [Medline](#)
143. Kang, J. W., Hong, J. M., and Lee, S. M. (2016) Melatonin enhances mitophagy and mitochondrial biogenesis in rats with carbon tetrachloride-induced liver fibrosis. *J. Pineal Res.* **60**, 383–393 [CrossRef](#) [Medline](#)
144. Yang, Y., Zhao, Z., Liu, Y., Kang, X., Zhang, H., and Meng, M. (2015) Suppression of oxidative stress and improvement of liver functions in mice by ursolic acid via LKB1-AMP-activated protein kinase signaling. *J. Gastroenterol. Hepatol.* **30**, 609–618 [CrossRef](#) [Medline](#)
145. Caligiuri, A., Bertolani, C., Guerra, C. T., Aleffi, S., Galastri, S., Trappolieri, M., Vizzutti, F., Gelmini, S., Laffi, G., Pinzani, M., and Marra, F. (2008) Adenosine monophosphate-activated protein kinase modulates the activated phenotype of hepatic stellate cells. *Hepatology* **47**, 668–676 [CrossRef](#) [Medline](#)

146. Kumar, P., Smith, T., Rahman, K., Thorn, N. E., and Anania, F. A. (2014) Adiponectin agonist ADP355 attenuates CCl<sub>4</sub>-induced liver fibrosis in mice. *PLoS ONE* **9**, e110405 [CrossRef](#) [Medline](#)
147. Zhai, X., Qiao, H., Guan, W., Li, Z., Cheng, Y., Jia, X., and Zhou, Y. (2015) Curcumin regulates peroxisome proliferator-activated receptor-gamma coactivator-1 $\alpha$  expression by AMPK pathway in hepatic stellate cells *in vitro*. *Eur. J. Pharmacol.* **746**, 56–62 [CrossRef](#) [Medline](#)
148. Zhang, W., Wu, R., Zhang, F., Xu, Y., Liu, B., Yang, Y., Zhou, H., Wang, L., Wan, K., Xiao, X., and Zhang, X. (2012) Thiazolidinediones improve hepatic fibrosis in rats with non-alcoholic steatohepatitis by activating the adenosine monophosphate-activated protein kinase signalling pathway. *Clin. Exp. Pharmacol. Physiol.* **39**, 1026–1033 [CrossRef](#) [Medline](#)
149. Dong, Z., Su, L., Esmaili, S., Iseli, T. J., Ramezani-Moghadam, M., Hu, L., Xu, A., George, J., and Wang, J. (2015) Adiponectin attenuates liver fibrosis by inducing nitric oxide production of hepatic stellate cells. *J. Mol. Med. (Berl.)* **93**, 1327–1339 [CrossRef](#) [Medline](#)
150. da Silva Morais, A., Abarca-Quinones, J., Guigas, B., Viollet, B., Stärkel, P., Horsmans, Y., and Leclercq, I. A. (2009) Development of hepatic fibrosis occurs normally in AMPK-deficient mice. *Clin. Sci. (Lond.)* **118**, 411–420 [CrossRef](#) [Medline](#)
151. Lim, J. Y., Oh, M. A., Kim, W. H., Sohn, H. Y., and Park, S. I. (2012) AMP-activated protein kinase inhibits TGF- $\beta$ -induced fibrogenic responses of hepatic stellate cells by targeting transcriptional coactivator p300. *J. Cell. Physiol.* **227**, 1081–1089 [CrossRef](#) [Medline](#)
152. Heinrichs, D., Knauel, M., Offermanns, C., Berres, M. L., Nellen, A., Leng, L., Schmitz, P., Bucala, R., Trautwein, C., Weber, C., Bernhagen, J., and Wasmuth, H. E. (2011) Macrophage migration inhibitory factor (MIF) exerts antifibrotic effects in experimental liver fibrosis via CD74. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 17444–17449 [CrossRef](#) [Medline](#)
153. Myers, R. W., Guan, H. P., Ehrhart, J., Petrov, A., Prahalada, S., Tozzo, E., Yang, X., Kurtz, M. M., Trujillo, M., Gonzalez Trotter, D., Feng, D., Xu, S., Eiermann, G., Holahan, M. A., Rubins, D., et al. (2017) Systemic pan-AMPK activator MK-8722 improves glucose homeostasis but induces cardiac hypertrophy. *Science* **357**, 507–511 [CrossRef](#) [Medline](#)